

A HAUSDORFF-YOUNG INEQUALITY FOR MEASURED GROUPOIDS

PATRICIA BOIVIN AND JEAN RENAULT

ABSTRACT. The classical Hausdorff-Young inequality for locally compact abelian groups states that, for $1 \leq p \leq 2$, the L^p -norm of a function dominates the L^q -norm of its Fourier transform, where $1/p + 1/q = 1$. By using the theory of non-commutative L^p -spaces and by reinterpreting the Fourier transform, R. Kunze (1958) [resp. M. Terp (1980)] extended this inequality to unimodular [resp. non-unimodular] groups. The analysis of the L^p -spaces of the von Neumann algebra of a measured groupoid provides a further extension of the Hausdorff-Young inequality to measured groupoids.

1. INTRODUCTION.

The classical Hausdorff-Young inequality for a locally compact abelian group G says that, for $1 \leq p \leq 2$ and for a compactly supported continuous function f on G , the L^p -norm of f dominates the L^q -norm of its Fourier transform $\hat{f}$, where q is the conjugate exponent ($1/p + 1/q = 1$):

$$\|\hat{f}\|_q \leq \|f\|_p.$$

The Fourier transform $\hat{f}$ is the function on the dual group $\hat{G}$ defined by

$$\hat{f}(\chi) = \int f(x) \overline{\chi(x)} dx.$$

The Haar measure of $\hat{G}$ is normalized so that the Fourier transform extends to an isometry from $L^2(G)$ onto $L^2(\hat{G})$. Thus, the inequality holds for $p = 2$. The proof is usually given as one of the first applications of the complex interpolation method, since the inequality also holds for $p = 1$.

This inequality has been generalized to non-abelian locally compact groups. The main difficulty is to give a meaning to the space $L^q(\hat{G})$ and to the Fourier transform $\mathcal{F}_p : L^p(G) \rightarrow L^q(\hat{G})$ when G is no longer abelian. The theory of operator algebras provides a solution. In the abelian case, the function $\hat{f}$ can be seen as the multiplication operator $M(\hat{f})$ on $L^2(\hat{G})$, i.e. $M(\hat{f})\xi = \hat{f}\xi$. Up to unitary equivalence, this is the convolution operator $L(f)$ on $L^2(G)$, i.e. $L(f)\xi = f*\xi$. This still makes sense when G is not abelian, provided one is willing to deal with unbounded operators. When f is continuous and compactly supported, $L(f)$ is a bounded operator on $L^2(G)$. The von Neumann algebra generated by these operators is the group von Neumann algebra $VN(G)$, which will play the rôle of the von Neumann algebra $L^\infty(\hat{G})$.

The case of unimodular groups was solved by R. Kunze in [10]. In that case, the von Neumann algebra $M = VN(G)$ has a trace τ and the non-commutative integration theory of I. Segal ([14]) is available. In fact, Kunze defines and studies

1991 *Mathematics Subject Classification.* Primary 46L55; Secondary 43A35, 43A07, 43A15, 43A22.

Key words and phrases. Von Neumann algebras. Non-commutative integration theory. Non-commutative L^p -spaces. Hausdorff-Young inequality.

non-commutative L^p -spaces $L^p(M, \tau)$ for a general von Neumann algebra M endowed with a trace τ . Its elements are unbounded operators on the Hilbert space $L^2(M, \tau)$ of the GNS construction affiliated with M . He then defines $L^q(\hat{G})$ as $L^q(VN(G), \tau)$. For $1 \leq p \leq 2$, the p -Fourier transform $\mathcal{F}_p : L^p(G) \rightarrow L^q(\hat{G})$ is given by $\mathcal{F}_p(f) = L(f)$ just as above.

One had to wait for the Tomita-Takesaki theory and its developments to tackle the non-unimodular case. Relying on the work of U. Haagerup ([7]) on one hand and of A. Connes ([4]) and M. Hilsuim ([8]) on the other on non-commutative L^p -spaces, M. Terp solved the general case of a locally compact group in a paper [17] unfortunately never published.

She also developed in [18], in the case of a weight instead of a state, the interpolation approach of H. Kosaki to non-commutative L^p -spaces, a framework particularly well-fitted to the generalization of the Hausdorff-Young inequality.

In the meantime, other Hausdorff-Young type inequalities were found. In particular, B. Russo gave in [13] the following Hausdorff-Young inequality for integral operators. Let X be a Borel space endowed with a σ -finite measure dx . Given $f \in L^2(X \times X)$, let $L(f)$ be the integral operator on $L^2(X)$ having f for kernel. Then, for $1 \leq p \leq 2$ and q conjugate exponent of p , the following inequality holds:

$$\|L(f)\|_q \leq \max(\|f\|_{p,q}, \|f^*\|_{p,q}),$$

where $\|L(f)\|_q$ is the q -Schatten norm of $L(f)$, $\|f\|_{p,q} = (\int (\int |f(x,y)|^p dx)^{q/p} dy)^{1/q}$ is the mixed norm introduced in [2] and $f^*(x,y) = f(y,x)$.

When $p = 1$, $q = \infty$; the corresponding norm is the usual operator norm and the inequality becomes

$$\|L(f)\| \leq \max(\sup_y \int |f(x,y)| dx, \sup_x \int |f(x,y)| dy)$$

which is a well-known consequence of the Cauchy-Schwarz inequality. When $p = 2$, $q = 2$; the corresponding norm is the Hilbert-Schmidt norm; both terms on the right hand side are equal by Fubini's theorem; as well known, we have the equality

$$\|L(f)\|_2 = (\int \int |f(x,y)|^2 dx dy)^{1/2}.$$

As in the classical case, the case $1 < p < 2$ is obtained by complex interpolation. Although the mixed norms $\|f\|_{p,q}$ are interpolation norms just as the usual L^p -norms are, one cannot conclude directly, due to the presence of a max of these norms.

The similarity between the group case and the case of integral operators is striking. It can still be made stronger by introducing the measured groupoid $G = X \times X$. Then the formulation is almost identical to the group case: the relevant von Neumann algebra is $M = VN(G)$ (it is the algebra of all bounded operators on $L^2(X)$ but acting by left multiplication on $L^2(G)$), $L(f)$ is the left convolution operator on $L^2(G)$ and the Schatten space S_q is the non-commutative L^q -space of M endowed with its natural trace. The main difference is that the usual L^p -norm of f has been replaced by the maximum of the mixed norms $\|f\|_{p,q}$ and $\|f^*\|_{p,q}$.

Let us fix our notation for groupoids. The unit space of the groupoid G will usually be denoted by $X = G^{(0)}$. We shall use greek letters like $\gamma, \dots$ for elements of G and roman letters $x, y, \dots$ for elements of $G^{(0)}$. The range and source maps will be denoted by $r, s : G \rightarrow G^{(0)}$. For $x \in G^{(0)}$, we define $G^x = r^{-1}(x)$ and $G_x = s^{-1}(x)$. The inverse of $\gamma \in G$ is denoted by γ^{-1} . A measured groupoid (see [5], [6]) consists of a Borel groupoid G , a measurable Haar system λ made of measures λ^x on G^x satisfying the left invariance and the measurability property, and a quasi-invariant measure μ on $G^{(0)}$. The quasi-invariance of μ is expressed

by the equivalence of the measures $\nu = \mu \circ \lambda$ and $\nu^{-1} = \mu \circ \tilde{\lambda}$, where $\tilde{\lambda}$ is the right invariant Haar system consisting of the measures $\lambda_x = (\lambda^x)^{-1}$ on G_x . By definition, the modular function is the Radon-Nikodym derivative $\delta = d\nu/d\nu^{-1}$. We say that the measured groupoid is unimodular if $\delta = 1$. To avoid technicalities, we assume that G is a second countable locally compact Hausdorff topological groupoid, that the measures are Radon measures, that λ is a continuous Haar system and that the modular function is continuous. Let us give some examples of measured groupoids: locally compact groups (with a chosen left Haar measure), measured spaces and more generally transformation groups (X, G) equipped with a quasi-invariant measure (with the appropriate assumptions to fit our hypotheses). As further examples: the above measured groupoid $X \times X$ and the holonomy groupoid of a foliation equipped with a Lebesgue measure. The mixed norms still make sense in this context. For f non-negative and measurable on G or in $C_c(G)$, one can define

$$\|f\|_{p,q} = \left(\int \left(\int |f|^p d\lambda^x \right)^{q/p} d\mu(x) \right)^{1/q}.$$

In the case of a group, $\|f\|_{p,q} = \|f\|_p$ while in the case of a space, $\|f\|_{p,q} = \|f\|_q$. On the other hand, we have the regular representation of G on $L^2(G)$ which defines the von Neumann algebra $M = VN(G)$ and provides the operator $L(f)$. Thus, we have all the ingredients to express a Hausdorff inequality for measured groupoids. As we shall see, it will have the following form

$$\|\mathcal{F}_p(f)\|_{L^q(M)} \leq \max(\|f\|_{p,q}, \|f^*\|_{p,q}),$$

where $1 \leq p \leq 2$, $\mathcal{F}_p(f) = \Delta^{1/2q} L(f) \Delta^{1/2q}$, Δ is the operator of multiplication by δ on $L^2(G)$ and $f^*(\gamma) = \overline{f(\gamma^{-1})}$.

2. NON-COMMUTATIVE L^p -SPACES FOR MEASURED GROUPOIDS.

Let (G, λ, μ) be a measured groupoid. As said above, we assume that G is a second countable locally compact Hausdorff topological groupoid, that the measures are Radon measures, that λ is a continuous Haar system and that the modular function δ is continuous. We denote by ν the measure $\nu = \mu \circ \lambda$ on G , by ν^{-1} its image under the inverse map and by ν_0 the symmetric measure $\nu_0 = \delta^{-1/2} \nu$.

It is convenient for our purpose to present the von Neumann algebra $M = VN(G)$ of the measured groupoid as the left algebra of a modular Hilbert algebra (see [15]). We endow the space $C_c(G)$ of compactly supported continuous functions on G with the convolution product

$$f * g(\gamma) = \int f(\gamma\eta) g(\eta^{-1}) d\lambda^{s(\gamma)}(\eta),$$

the involution

$$f^*(\gamma) = \overline{f(\gamma^{-1})}$$

the inner product

$$(f|g) = \int f(\gamma) \overline{g(\gamma)} d\nu^{-1}(\gamma)$$

and the complex one-parameter group $\Delta(z)$ of automorphisms of $C_c(G)$

$$\Delta(z)f = \delta^z f.$$

The axioms (I) to (VIII) of [15, Definition 2.1] are readily verified. The left and right convolution operators $L(f)$ and $R(g)$ are respectively defined by

$$L(f)\xi = f * \xi \quad R(g)\eta = \eta * g.$$

They extend to bounded linear operators, still denoted by $L(f)$ and $R(g)$, on the completed Hilbert space $H = L^2(G, \nu^{-1})$. The map L is a representation of the

$*$ -algebra $C_c(G)$. We define $M = VN(G)$ as the von Neumann algebra generated by $\{L(f), f \in C_c(G)\}$. According to the general theory, the von Neumann algebra generated by $\{R(g), g \in C_c(G)\}$ is the commutant M' of M . The polar decomposition $S = J\Delta^{1/2}$ of the closure S of the operator $\xi \mapsto \xi^*$, viewed as an unbounded antilinear operator on H provides the ingredients of the Tomita-Takesaki theory. The modular operator Δ is the operator of multiplication by δ ; it is self-adjoint positive; its domain $Dom(\Delta)$ is the subspace of the ξ 's in H such that $\delta\xi \in H$; it contains $C_c(G)$ as an essential domain. The modular involution J is given by $J\xi = \delta^{1/2}\xi^*$. The canonical weight φ_0 on M is defined by

$$\varphi_0(L(f)^*L(g)) = (g|f) \quad \text{for } f, g \in C_c(G).$$

Another useful formula is

$$\varphi_0(L(f)) = \int f|_{G^{(0)}} d\mu \quad \text{for } f \in C_c(G).$$

The modular group of the weight φ_0 is given by $\sigma_t(T) = \Delta^{it}T\Delta^{-it}$ for $T \in M$.

The spatial theory (see [4, 8]) uses the dual weight ψ_0 on M' , defined by $\psi_0(T) = \varphi_0(JTJ)$. We have

$$\psi_0(R(Jf)^*R(Jg)) = (g|f) \quad \text{for } f, g \in C_c(G).$$

In our context, it can be expressed as follows. A vector $\xi \in H$ is called left bounded if the operator $L(\xi)$ defined by $L(\xi)f = \xi * f$ for $f \in C_c(G)$ is bounded. The space of left bounded operators is denoted by $\mathcal{A}_l$. One associates to each normal semi-finite weight φ on M a self-adjoint positive operator T on H such that

$$\|T^{1/2}\xi\|^2 = \varphi(L(\xi)L(\xi)^*)$$

for all $\xi \in \mathcal{A}_l$. The operator T is called the spatial Radon-Nikodym derivative of the weight φ and is denoted by $\frac{d\varphi}{d\psi_0}$. It is called integrable if φ is a state.

One then defines its integral as $\int T d\psi_0 = \varphi(1)$. The correspondence $\varphi \mapsto \frac{d\varphi}{d\psi_0}$ extends to $\varphi \in M_*$. The space $L^1(M, \psi_0)$ is defined as its image. On the other hand, if $\varphi \in M_*$, $f \mapsto \varphi(L(f))$ is a Radon measure on $C_c(G)$ which is absolutely continuous with respect to $\nu = \mu \circ \lambda$. We denote by $\frac{d\varphi}{d\nu_0}$ its usual Radon-Nikodym derivative with respect to the symmetric measure ν_0 . Explicitly, if $\varphi(T) = (T\xi|\eta)$ with $\xi, \eta \in H$, one finds that $\frac{d\varphi}{d\nu_0}$ is the coefficient

$$(\xi', \eta')(\gamma) = (L(\gamma)\xi' \circ s(\gamma)|\eta' \circ r(\gamma))$$

of the left regular representation of G on the field of Hilbert spaces $x \mapsto L^2(G^x, \lambda^x)$, where $\xi' = \delta^{-1/2}\xi$ and $\eta' = \delta^{-1/2}\eta$. We define the space $\mathcal{L}^1(\hat{G})$ as the image of the map $\varphi \in M_* \mapsto \frac{d\varphi}{d\nu_0}$. The three spaces M_* , $L^1(M, \psi_0)$ and $\mathcal{L}^1(\hat{G})$ are isomorphic as linear spaces, and also as Banach spaces with the appropriate norms. When $\varphi(T) = (T\xi|\xi)$ with $\xi \in C_c(G)$, the spatial Radon-Nikodym derivative $T = \frac{d\varphi}{d\psi_0}$ is the closure of the operator $\Delta^{1/2}L(\tilde{F})\Delta^{1/2}$ defined on $C_c(G)$, where $F = \frac{d\varphi}{d\nu_0}$ and $\tilde{F}(\gamma) = F(\gamma^{-1})$. When G is a locally compact group, $\mathcal{L}^1(\hat{G})$ is the usual Fourier algebra $A(G)$.

The Hilbert space $H = L^2(G, \nu^{-1})$ admits the direct integral decomposition $H = \int^\oplus H_x d\mu_x$, where $H_x = L^2(G_x, \lambda_x)$. A decomposable operator $T = \int^\oplus T_x d\mu_x$

is called α -homogeneous if for a.e. $\gamma \in G$,

$$R(\gamma)T_{s(\gamma)} \subset \delta^{-\alpha}(\gamma)T_{r(\gamma)}R(\gamma),$$

where $R(\gamma)$ is the isometry $H_{s(\gamma)} \rightarrow H_{r(\gamma)}$ given by $R(\gamma)\xi(\gamma') = \xi(\gamma'\gamma)$. The elements of M (and more generally the operators affiliated with M) are 0-homogeneous. The operator Δ as well as the elements of $L^1(M, \psi_0)$ are (-1) -homogeneous. For $p \in [0, \infty]$, $L^p(M, \psi_0)$ is the set of closed densely defined $(-1/p)$ -homogeneous operators H such that $|T|^p$ belongs to $L^1(M, \psi_0)$. The p -norm of $T \in L^p(M, \psi_0)$ is defined as $\|T\|_p = (\int |T|^p d\psi_0)^{1/p}$. For $p = \infty$, we define $L^\infty(M, \psi_0) = M$.

The well-known isomorphism of $L^2(X \times X)$ onto the space of Hilbert-Schmidt operators on $L^2(X)$ which associates to f the integral operator $L(f)$ with kernel f and the Fourier-Plancherel transform of $L^2(G)$ onto $L^2(\hat{G})$, where G is a locally compact abelian group can be viewed as particular cases of the following general result for measured groupoids, which is proved in [3].

Theorem 2.1. *Let (G, λ, μ) be a measured groupoid. There exists a unique isometry $\mathcal{F}_2 : L^2(G, \nu_0) \rightarrow L^2(M, \psi_0)$ such that, for $f \in C_c(G)$, $\mathcal{F}_2(f)$ is the closure of the operator $\Delta^{1/4}L(f)\Delta^{1/4}$ defined on $C_c(G)$.*

In [18], M. Terp defines the non-commutative L^p -spaces by the complex interpolation method in a framework perfectly well-fitted to ours. The data consists of a von Neumann algebra M and a (normal, faithful and semi-finite) weight φ . She defines the “intersection” L of M and its predual M_* and then embeds M and M_* into L^* , turning (M, M_*) into a compatible pair of Banach spaces. She also defines (Section 2.3) for all $p \in [1, \infty]$ a linear, norm-decreasing, injective map $\mu_p : L \rightarrow L^p(M, \psi)$ which has dense range for $p < \infty$ and where ψ is the dual weight of M' in the GNS representation of (M, φ) . We apply this to $(M = VN(G), \varphi = \varphi_0)$. We observe that $C_c(G) * C_c(G)$ is contained in L and that for $f \in C_c(G) * C_c(G)$, $\mu_p(f)$ is the closure of the operator $\Delta^{1/2p}L(f)\Delta^{1/2p}$ defined on $C_c(G)$. For $p = 1$, $f \mapsto \mu_1(\tilde{f})$ extends to the isomorphism $\mathcal{L}^1(\hat{G}) \rightarrow L^1(M, \psi_0)$ described earlier. For $p = 2$, $f \mapsto \mu_2(f)$ extends to the above Fourier-Plancherel isometry $\mathcal{F}_2 : L^2(G, \nu_0) \rightarrow L^2(M, \psi_0)$. For $p = \infty$, $\mu_\infty = L$ is the left regular representation.

3. A HAUSDORFF-YOUNG INEQUALITY.

We use the same notation as in the previous section. Because of our assumptions, Δ as well as Δ^z , where $z \in \mathbf{C}$, admit $C_c(G)$ as an essential domain.

Lemma 3.1. *Let $z \in \mathbf{C}$ and $f \in C_c(G)$. We have the commutation rule*

$$L(f)\Delta^{-z} \subset \Delta^{-z}L(\delta^z f).$$

Proof. We have seen that Δ^z implements the automorphism $\Delta(z)$ of the modular Hilbert algebra $C_c(G)$. Therefore, $\Delta^z L(f)\Delta^{-z} \subset L(\delta^z f)$. $\square$

Since Δ^z as well as $L(f)$ leave $C_c(G)$ invariant, $C_c(G)$ is contained in the domain of $\Delta^z L(f)\Delta^z$. Therefore $\Delta^z L(f)\Delta^z$ admits an adjoint; moreover the equality

$$(\Delta^z L(f)\Delta^z \xi | \eta) = (\xi | \Delta^{\bar{z}} L(f^*) \Delta^{\bar{z}} \eta)$$

for $\xi, \eta \in C_c(G)$ shows that this adjoint is an extension of $\Delta^{\bar{z}} L(f^*) \Delta^{\bar{z}}$. In particular, $\Delta^z L(f)\Delta^z$ is closable.

Definition 3.1. Given $\alpha \in \mathbf{C}$ and $\beta = (1 - 2\alpha)^{-1}$, we define the β -Fourier transform $\mathcal{F}_\beta(f)$ of $f \in C_c(G)$ as the closure of the operator $\Delta^\alpha L(f)\Delta^\alpha$ defined on $C_c(G)$.

We have met particular cases of this definition: when f belongs to $C_c(G) * C_c(G)$ and $\alpha = 1/2q$, where $q \in [1, \infty]$, then $\beta = p$ is the conjugate exponent of q and $\mathcal{F}_p(f) = \mu_q(f)$, where μ_q is Terqs symmetric embedding recalled earlier. In that case, $\mathcal{F}_p(f)$ belongs to $L^q(M, \psi_0)$. On the other hand, $\mathcal{F}_2(f)$ belongs to $L^2(M, \psi_0)$ and $\mathcal{F}_1(f) = L(f)$ belongs to $L^\infty(M, \psi_0) = M$ for all $f \in C_c(G)$.

Our goal is to show that, for $1 \leq p \leq 2$, and $f \in C_c(G)$, $\mathcal{F}_p(f)$ belongs to $L^q(M, \psi_0)$ where q is the conjugate exponent of p . and satisfies the following Hausdorff-Young inequality:

$$\|\mathcal{F}_p(f)\|_q \leq \max(\|f\|_{p,q}, \|f^*\|_{p,q})$$

where $\|f\|_{p,q}$ denotes the following mixed norm, which generalizes the mixed norms of [2].

Definition 3.2. Given $p, q \in [1, \infty]$ and $f \in C_c(G)$, the mixed norm $\|f\|_{p,q}$ of f is defined as

$$\|f\|_{p,q} = \left(\int \left(\int |f|^p d\lambda^x \right)^{q/p} d\mu(x) \right)^{1/q}$$

with the obvious modifications when p or q is infinite.

We first verify that this inequality holds when $p = 1$ or $p = 2$. Indeed, for $p = 1$, this is the well-known inequality

$$\|L(f)\| \leq \max(\|f\|_{1,\infty}, \|f^*\|_{1,\infty})$$

which is obtained by the Cauchy-Schwarz inequality. For $p = 2$, we have

$$\|\mathcal{F}_2(f)\|_2^2 = \int |f|^2 d\nu_0 \leq \left(\int |f|^2 d\nu \right)^{1/2} \left(\int |f|^2 d\nu^{-1} \right)^{1/2} = \|f\|_{2,2} \|f^*\|_{2,2}$$

again by Cauchy-Schwarz. We want to deduce the general case $1 \leq p \leq 2$ from these two extremal cases by interpolation. Although the norm $\|\cdot\|_{p,q}$ interpolates between 1 and 2 (in fact between 1 and ∞), we cannot interpolate directly because our formula involves the maximum of two mixed norms. We proceed exactly as in [17]. We shall only give the necessary modifications. It relies on the following characterization of $L^p(M, \psi_0)$.

Proposition 3.2. [17, Proposition 2.3] *Let $p \in [1, \infty]$ and define q by $1/p + 1/q = 1$. Let T be a closed densely defined $(-1/p)$ -homogeneous operator on H . Then, the following conditions are equivalent:*

- (i) $T \in L^p(M, \psi_0)$,
- (ii) *there exists a constant $C \geq 0$ such that*

$$\forall S \in L^q(M, \psi_0), \forall \xi \in \mathcal{A}_l \cap \text{Dom}(T), \forall \eta \in \mathcal{A}_l \cap \text{Dom}(S) :$$

$$|(T\xi, S\eta)| \leq C \|S\|_q \|L(\xi)\| \|L(\eta)\|.$$

If $T \in L^p(M, \psi_0)$, then $\|T\|_p$ is the smallest C satisfying (ii).

In the sequel, we assume that $1 < p < 2$, we define q by $1/p + 1/q = 1$ and we let $f \in C_c(G)$ such that $\max(\|f\|_{p,q}, \|f^*\|_{p,q}) \leq 1$. We let $M(x, y) = \max(\|f^x\|_p, \|f^y\|_p)$. We choose $\epsilon > 0$ such that $\epsilon^{(q-p)} \|f\|_p^p \leq 1$ and we set $M_\epsilon(x, y) = \max(M(x, y), \epsilon)$. We define for $z \in \mathbf{C}$ and $\gamma \in G$

$$\begin{aligned} f_z(\gamma) &= \text{sgn} f(\gamma) |f(\gamma)|^{p_z} M_\epsilon(r(\gamma), s(\gamma))^{q-z(p+q)} \quad \text{if } f(\gamma) \neq 0 \\ f_z(\gamma) &= 0 \quad \text{if } f(\gamma) = 0. \end{aligned}$$

Then f_z belongs to $C_c(G)$. We denote by Ω the open strip $\{z \in \mathbf{C} : 1/2 \leq \Re z \leq 1\}$. We have the following lemma which is proved exactly as [17, Lemma 4.1].

Lemma 3.3. *Let $p \in [1, 2]$, $\Omega \subset \mathbf{C}$, $f, f_z \in C_c(G)$ be as above. Let $\xi \in \mathcal{A}_l$. Then*

(i) for each $z \in \overline{\Omega}$, the function

$$\xi_z = \mathcal{F}_{1/z}(f_z)\xi = \Delta^{\frac{1-z}{2}}L(f_z)\Delta^{\frac{1-z}{2}}\xi$$

is defined and belongs to H ;

(ii) the function $z \mapsto \xi_z$ with values in H is bounded on $\overline{\Omega}$;

(iii) for each $\eta \in H$, the scalar function $z \mapsto (\xi_z, \eta)$ is continuous on $\overline{\Omega}$ and analytic on Ω .

Proof. The proof given in [17, Lemma 4.1] applies with little change. To prove (i) and (ii), one introduces $\eta \in C_c(G)$ and considers the integral

$$H_\eta(z) = \int \xi(\gamma) \overline{\mathcal{F}_{1/\bar{z}}(f_z^*)\eta(\gamma)} d\nu^{-1}(\gamma).$$

By the use of Fubini's theorem, one checks that this is well defined and that

$$H_\eta(z) = \int \xi_z(\gamma) \overline{\eta(\gamma)} d\nu^{-1}(\gamma).$$

Then, one proves that there exists $C \geq 0$ such that $|H_\eta(z)| \leq \|\eta\|_2$. To do that, one applies the Phragmen-Lindelöf principle. The function $z \rightarrow H_\eta(z)$ is bounded continuous on $\overline{\Omega}$ and analytic on Ω . It suffices to estimate it on the lines $\Re z = 1$ and $\Re z = 1/2$.

Suppose that $z = 1 + it$, where t is real. Then,

$$\xi_z = \Delta^{-it/2}L(f_z)\Delta^{-it/2}\xi$$

belongs to H and we have

$$|H_\eta(z)| \leq \|L(f_z)\| \|\xi\|_2 \|\eta\|_2.$$

In order to estimate the operator norm of $L(f_z)$, we use the inequality

$$\|L(f_z)\| \leq \max(\|f_z\|_{1,\infty}, \|f_z^*\|_{1,\infty}).$$

Let us fix $x \in G^{(0)}$ and estimate

$$\int |f_z| d\lambda^x = \int |f(\gamma)|^p M_\epsilon(x, s(\gamma))^{-p} d\lambda^x(\gamma).$$

We write G^x as the disjoint union $A \cup B$, where $A = \{\gamma \in G^x : M(x, s(\gamma)) \leq \epsilon\}$ and $B = G^x \setminus A$. If $\gamma \in A$, $M_\epsilon(x, s(\gamma)) = M(x, s(\gamma)) \geq \|f^x\|_p$. Therefore

$$\int_A |f_z| d\lambda^x \leq \|f^x\|_p^{-p} \int_A |f(\gamma)|^p d\lambda^x(\gamma) \leq 1.$$

On the other hand, if B is non-empty, then $\|f^x\|_p < \epsilon$ and for all $\gamma \in B$, $M_\epsilon(x, s(\gamma)) = \epsilon$. Therefore,

$$\int_B |f_z| d\lambda^x \leq \epsilon^{-p} \int_B |f(\gamma)|^p d\lambda^x(\gamma) \leq \epsilon^{-p} \epsilon^p \leq 1.$$

This shows that $\|f_z\|_{1,\infty} \leq 2$. One shows similarly that $\|f_z^*\|_{1,\infty} \leq 2$. This gives $\|L(f_z)\| \leq 2$ and

$$|H_\eta(z)| \leq 2\|\xi\|_2\|\eta\|_2.$$

Suppose next that $z = \frac{1}{2} + it$, where t is real. Then,

$$\xi_z = \Delta^{-it}\mathcal{F}_2(\delta^{it/2}f_z)\xi.$$

This also belongs to H because the domain of operators in $L^2(M, \psi_0)$ contains $\mathcal{A}_l$.

We deduce the inequality

$$|H_\eta(z)| \leq \|\mathcal{F}_2(\delta^{it/2}f_z)\xi\|_2\|\eta\|_2.$$

As a corollary of Proposition 3.2, we have the inequality

$$\|\mathcal{F}_2(\delta^{it/2}f_z)\xi\|_2 \leq \|\mathcal{F}_2(\delta^{it/2}f_z)\|_2 \|L(\xi)\|.$$

We use the Plancherel Theorem 2.1 to evaluate the norm

$$\|\mathcal{F}_2(\delta^{it/2} f_z)\|_2 = \left(\int |f_z|^2 d\nu_0 \right)^{1/2} \leq \left(\int |f_z|^2 d\nu \int |f_z|^2 d\nu^{-1} \right)^{1/4}.$$

Let us estimate the integral

$$\int |f_z|^2 d\nu = \int |f(\gamma)|^p M_\epsilon(r(\gamma), s(\gamma))^{(q-p)} d(\mu \circ \lambda)(\gamma).$$

We write the domain of summation as the union of three parts

$$\begin{aligned} A &= \{\gamma : \|f^{r(\gamma)}\|_p \geq \max(\|f_{s(\gamma)}\|_p, \epsilon)\}, \\ B &= \{\gamma : \|f_{s(\gamma)}\|_p \geq \max(\|f^{r(\gamma)}\|_p, \epsilon)\} \end{aligned}$$

and

$$C = \{\gamma : M(r(\gamma), s(\gamma)) \leq \epsilon\}.$$

If $\gamma \in A$, $M_\epsilon(r(\gamma), s(\gamma)) = \|f^{r(\gamma)}\|_p$, therefore

$$\int_A |f_z|^2 d(\mu \circ \lambda) \leq \int \left(\int |f(\gamma)|^p d\lambda^x(\gamma) \right) \|f^x\|_{\bar{p}}^{-p} \|f^x\|_p^q d\mu(x) = \|f\|_{p,q}^q \leq 1.$$

Similarly,

$$\int_B |f_z|^2 d(\mu \circ \lambda) \leq \|f^*\|_{p,q}^q \leq 1.$$

Finally, if $\gamma \in C$, $M_\epsilon(r(\gamma), s(\gamma)) = \epsilon$ and

$$\int_C |f_z|^2 d(\mu \circ \lambda) = \epsilon^{(q-p)} \int_R |f|^p d(\mu \circ \lambda) \leq \epsilon^{(q-p)} \|f\|_p^p.$$

Because of our choice of $\epsilon > 0$, this is majorized by 1. The whole integral is majorized by 3. One shows in the same fashion that the integral $\int |f_z|^2 d\nu^{-1}$ is also majorized by 3. Therefore, $\|\mathcal{F}_2(\delta^{it/2} f_z)\|_2 \leq \sqrt{3} \leq 2$. This gives

$$|H_\eta(z)| \leq 2\|L(\xi)\| \|\eta\|_2$$

on the line $\Re z = 1/2$.

This proves the desired inequality with $C = \max(\|\xi\|_2, \|L(\xi)\|)$ and (i) and (ii). We have already proved (iii) for $\eta \in C_c(G)$. The general case follows by approximation. $\square$

The proof of the following lemma, which is exactly [17, Lemma 4.2], does not require any modification.

Lemma 3.4. *Let $p \in [1, 2]$ and let q be defined by $1/p + 1/q = 1$. Let $f \in C_c(G)$ be as above and let $S \in L^p(M, \psi_0)$. Then, for all $\xi \in \mathcal{A}_l$ and $\eta \in \mathcal{A}_l \cap \text{Dom}(S)$, we have:*

$$|(\mathcal{F}_p(f)\xi, S\eta)| \leq 2\|S\|_p \|L(\xi)\| \|L(\eta)\|.$$

Note that $\xi \in \text{Dom}(\mathcal{F}_p(f))$ by Lemma 3.3.

Proof. We just recall how the proof goes. We assume $\|S\|_p \leq 1$. By [17, Lemma 2.5], it suffices to consider the case $\eta \in \mathcal{A}_l \cap \text{Dom}(|S|^p)$. Let $S = U|S|$ be the polar decomposition of S . For $z \in \overline{\Omega}$, one defines

$$\eta_z = U|S|^{pz}\eta.$$

Then the function $z \mapsto \eta_z \in H$ is bounded continuous on $\overline{\Omega}$ and analytic on Ω . We then define $H(z) = (\xi_z, \eta_z)$. The scalar function $z \mapsto H(z)$ satisfies the same properties. One proves the estimate

$$|H(z)| \leq 2\|L(\xi)\| \|L(\eta)\|$$

for $z \in \Omega$ by checking it on the lines $\Re z = 1$ and $\Re z = 1/2$ as in the previous lemma.

For $z = 1 + it$, where t is real, one has

$$\begin{aligned} |H(z)| &= (L(\delta^{it/2} f_z) \xi, \Delta^{it} U |S|^{it} |S|^p \eta) \\ &\leq \|L(\delta^{it/2} f_z)\| \|L(\xi)\| \|L(\eta)\|. \end{aligned}$$

by applying Proposition 3.2 with $p = \infty$. Then one uses the estimate

$$\|L(\delta^{it/2} f_z)\| \leq 2$$

established in the previous lemma.

For $z = \frac{1}{2} + it$, where t is real, one simply uses Cauchy-Schwarz' inequality

$$|H(z)| \leq \|\xi_z\|_2 \|\eta_{\bar{z}}\|_2.$$

We have established earlier the estimate $\|\xi_z\|_2 \leq 2\|L(\xi)\|$. On the other hand,

$$\|\eta_{\bar{z}}\|_2 = \|U |S|^{p/2} |S|^{-pit} \eta\|_2$$

can be also estimated by Proposition 3.2 with $p = 2$:

$$\|U |S|^{p/2} |S|^{-pit} \eta\|_2 \leq \|U |S|^{p/2}\|_2 \|L(|S|^{-pit} \eta)\| \leq \|L(\eta)\|.$$

In particular, we have

$$|H(1/p)| = |(\mathcal{F}_p(f) \xi, S \eta)| \leq 2\|L(\xi)\| \|L(\eta)\|.$$

□

One deduces from Proposition 3.2 that for $f \in C_c(G)$, $\mathcal{F}_p(f)$ belongs to $L^q(M, \psi_0)$ for all $p \in [1, 2]$ where $1/p + 1/q = 1$ and

$$\|\mathcal{F}_p(f)\|_{L^q(M)} \leq 2 \max(\|f\|_{p,q}, \|f^*\|_{p,q}).$$

This inequality can be improved.

Theorem 3.5. (*Hausdorff-Young*) *Let $p \in [1, 2]$ and $f \in C_c(G)$. Then $\mathcal{F}_p(f)$ belongs to $L^q(M, \psi_0)$ where $1/p + 1/q = 1$ and*

$$\|\mathcal{F}_p(f)\|_{L^q(M)} \leq \max(\|f\|_{p,q}, \|f^*\|_{p,q})$$

where $1/p + 1/q = 1$.

Proof. We first observe that the mixed norms as well as the L^p norms behave well with respect to products of groupoids. Explicitly, let (G_i, λ_i, μ_i) , $i = 1, 2$ be two measured groupoids and let $(G = G_1 \times G_2, \lambda = \lambda_1 \otimes \lambda_2, \mu = \mu_1 \otimes \mu_2)$ be their product. Given $1 \leq p, q \leq \infty$ and $f_i \in C_c(G)$, we have

- $\|f_1 \otimes f_2\|_{p,q} = \|f_1\|_{p,q} \|f_2\|_{p,q}$;
- $\mathcal{F}_p(f_1 \otimes f_2) = \mathcal{F}_p^{(1)}(f_1) \otimes \mathcal{F}_p^{(2)}(f_2)$;
- for $T_i \in L^q(M^{(i)}, \psi_0^{(i)})$, $T = T_1 \otimes T_2$ belongs to $L^q(M, \psi_0)$ and $\|T\|_{L^q(M)} = \|T_1\|_{L^q(M^{(1)})} \|T_2\|_{L^q(M^{(2)})}$ where the indices 1, 2 refer to the corresponding groupoids. The Hilbert space, the von Neumann algebra and the canonical weights of the product groupoid are also tensor products: $H = H^{(1)} \otimes H^{(2)}$, $M = M^{(1)} \otimes M^{(2)}$, $\varphi = \varphi^{(1)} \otimes \varphi^{(2)}$ and $\psi = \psi^{(1)} \otimes \psi^{(2)}$.

Given $f \in C_c(G)$, we form

$$F = (f^* \otimes f) \otimes (f^* \otimes f) \otimes \dots (f^* \otimes f)$$

where $f^* \otimes f$ is repeated n times. It belongs to $C_c(G^{2n})$. According to the above $\mathcal{F}_p(F)$ belongs to $L^q(M^{\otimes 2n}, \psi_0^{\otimes 2n})$ and

$$\|\mathcal{F}_p(F)\|_{L^q(M)} \leq 2 \max(\|F\|_{p,q}, \|F^*\|_{p,q}).$$

Since $\|F\|_{p,q} = \|F^*\|_{p,q} = \|f\|_{p,q}^n \|f^*\|_{p,q}^n$ and $\|\mathcal{F}_p(F)\|_{L^q(M^{2n})} = \|\mathcal{F}_p(f)\|_{L^q(M)}$, this gives

$$\|\mathcal{F}_p(f)\|_{L^q(M)} \leq 2^{1/2n} \|f\|_{p,q}^{1/2} \|f^*\|_{p,q}^{1/2}.$$

Passing to the limit, one gets:

$$\|\mathcal{F}_p(f)\|_{L^q(M)} \leq \|f\|_{p,q}^{1/2} \|f^*\|_{p,q}^{1/2} \leq \max(\|f\|_{p,q}, \|f^*\|_{p,q}).$$

□

REFERENCES

- [1] C. Anantharaman et J. Renault, *Amenable groupoids*, Contemp. Math., 282, Amer. Math. Soc., Providence, RI, 2001.
- [2] A. Benadek et R. Panzone, *The spaces L^p with mixed norm*, Duke Math. J., **28** (1961), 301-324.
- [3] P. Boivin, *Espaces L^p non-commutatifs de l'algèbre de von Neumann d'un groupoïde mesuré*, Thèse, Université d'Orléans, 2007.
- [4] A. Connes, *On the spatial theory of von Neumann algebras*, J. Func. Anal., **35** (1980), 153-164.
- [5] P. Hahn, *Haar measure for measure groupoids*, Trans. Amer. Math. Soc., **242** (1978).
- [6] P. Hahn, *Regular Representations of measure groupoids*, Trans. Amer. Math. Soc., **242** (1978).
- [7] U. Haagerup, *L^p -spaces associated with an arbitrary von Neumann algebra*, Algèbres d'opérateurs et leurs applications en physique mathématique (Proc. Colloq., Marseille, 1977), Colloq. Internat. CNRS, **274** (1979), 175-184.
- [8] M. Hilsu, *Les espaces L^p d'une algèbre de von Neumann dé finies par la dé rivée spatiale*, J. Funct. Anal., **40** (1981), 151-169.
- [9] H. Kosaki, *Applications of the Complex Interpolation Method to a von Neumann Algebra : Non-commutative L^p -Spaces*, J. of Funct. Anal., **56** (1984), 29-78.
- [10] R. A. Kunze, *L_p Fourier Transforms on locally compact unimodular groups*, Trans. Amer. Math. Soc, **89** (1958), 519-540.
- [11] P. S. Muhly, J. N. Renault and D. P. Williams, *Equivalence and isomorphism for groupoid C^* -algebras*, J. Operator Theory, **17** (1987), 3-22.
- [12] J. Renault: *A groupoid approach to C^* -algebras*, Lecture Notes in Mathematics, Vol. **793** Springer-Verlag Berlin, Heidelberg, New York (1980).
- [13] B. Russo, *On the Hausdorff-Young theorem for integral operators*, Pacific J. Math., **68** (1977), 241-253.
- [14] I. E. Segal, *A non-commutative extension of abstract integration*, Ann. of Math., **57** (1953), n°3.
- [15] M. Takesaki, *Tomita's Theory of Modular Hilbert Algebras and its Applications*, Lecture Notes in Math., **128**, Springer, 1970.
- [16] S. Strătilă and L. Zsidő, *Lectures on von Neumann Algebras*, Abacus Press, Kent, 1979.
- [17] M. Terp, *L^p Fourier Transformation on non-unimodular locally compact groups* Kobenhavns Universitet Matematisk Institut, Preprint, 1980.
- [18] M. Terp, *Interpolation spaces between a von Neumann algebra and its predual*, J. Operator Theory, **8** (1982), 327-360.

DÉPARTEMENT DE MATHÉMATIQUES, UNIVERSITÉ D'ORLÉANS, 45067 ORLÉANS, FRANCE
E-mail address: Jean.Renault@univ-orleans.fr, boivin.patricia@wanadoo.fr